

BREXIL MIX

BREXIL MIX

F1873

BREXIL MIX

BREXIL MIX

SOLUBLE MICRONUTRIENT AND SECONDARY NUTRIENT MICROGRANULES

GUARANTEED ANALYSIS

Magnesium (Mg)	3.20 % water soluble Magnesium (Mg)	3.20 %
Boron (B)		0.96 %
Copper (Cu)	0.64 % water soluble Copper (Cu)	0.64 %
Manganese (Mn)	0.56 % water soluble Manganese (Mn)	0.56 %
Molybdenum (Mo)		0.80 %
Zinc (Zn)	4.60 % water soluble Zinc (Zn)	4.60 %

Derived From: Copper Ligninsulfonate, Zinc Ligninsulfonate, Manganese Ligninsulfonate, Magnesium Ligninsulfonate, Boron Ethanolamine and Sodium Molybdate.

GENERAL INFORMATION

BREXIL MIX (0-0-0) is a micronutrient and secondary nutrient based product designed to treat and prevent micronutrient and secondary nutrient deficiencies in all crops. The ligninsulfonate (LSA) derived ligninpolycarboxylic acid (LPCA) in **BREXIL MIX** allows for quick absorption of the nutrients into the plant.

BREXIL MIX is a microgranule product that is completely water soluble. **BREXIL MIX** is only recommended as a supplement to a well balanced fertilizer program.

WARNING

This product contains Boron (B) and may be harmful to certain crops. Use only according to manufacturer's directions.

This product contains Molybdenum (Mo). Crops high in molybdenum can be toxic to ruminant animals. Use only according to manufacturer's directions.

A compatibility jar test must be performed before mixing **BREXIL MIX** with products containing Calcium and/or phosphate fertilizer.

DO NOT treat peaches, apricots and kiwis after fruit has reached half of expected size. DO NOT apply via fertigation.

Some crops may be injured by the application of boron and molybdenum. Contact your county agent or field representative before making applications.

FIELD APPLICATION INSTRUCTIONS, FOLIAR RATES		
CROP	RATE	Application Timing
Vegetable crops Including but not limited to: pepper, eggplant, zucchini, melon, water melon, cucumber, tomato, cole crops, leafy vegetables	Maintenance: 16 oz/acre Deficiency Symptoms: 24-32 oz/acre	Apply every 14-21 days intervals until deficiency symptoms disappear
Fruit crops Including but not limited to: apples, cherries, citrus, pear, stone fruit, kiwi, raisin, table and wine grapes, berry crops such as blueberries, cranberries and strawberries, etc.	Maintenance: 16 oz/acre Deficiency Symptoms: Apples and pears: 32-40 oz/acre; Stone fruit: 28-40 oz/acre; Citrus and grapes: 32-40 oz/acre; Berries: 24-32 oz/acre	Apply every 14-21 days intervals until deficiency symptoms disappear
Nut crops Including but not limited to: almond, pecan, pistachio, walnut,	Maintenance: 16 oz/acre Deficiency Symptoms: 28-40 oz/acre	Apply every 14-21 days intervals until deficiency symptoms disappear
Field crops Including but not limited to: corn, cotton, peanuts, potatoes, rice, soybeans, sorghum, sugar beets, sugar cane, wheat, etc.	Maintenance: 8 oz/acre Deficiency Symptoms: 16 oz/acre	Apply every 14-21 days intervals until deficiency symptoms disappear
Forage crops Including but not limited to: alfalfa, timothy, fescue, etc.	Maintenance: 8 oz/acre Deficiency Symptoms: 16 oz/acre	Apply every 14-21 days intervals until deficiency symptoms disappear
Ornamentals	Maintenance: 16 oz/acre Deficiency Symptoms: 24-32 oz/acre	Apply every 14-21 days intervals until deficiency symptoms disappear
Turf	1-3 oz/1000 sq ft	Apply every 14-21 days intervals until deficiency symptoms disappear
Greenhouse	Not more than 10 oz/25 Gallons of water	
Low volume spray (25 GPA)	Not more than 16 oz/acre	

Spray Volumes:

The standard spray volume on horticultural crops is 50 gallons per acre.

For spray volumes 5-20 GPA, use no more than 1lb/acre.

For spray volumes less than 5 GPA, use no more than ½ lb/acre.

In green house applications, do NOT apply more than 10 oz/25 gallons of spray.

5
Lb



NET WEIGHT
5Lb (2.3 kg)

DIRECTIONS FOR MIXING

BREXIL MIX is a soluble product to be mixed with water.

Select a water volume and corresponding rate of **BREXIL MIX** necessary to thoroughly spray/mist all fruit and foliage surfaces.

Gradually add the recommended amount of product to a half-filled spray tank and mix, then add the balance of required water while continuing to agitate the solution.

STORAGE INSTRUCTION

Keep the product sheltered from the sun and in a dry warehouse. Once open, completely use the contents of the bag.

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid breathing spray mist. May cause irritation of nose, throat, or skin. Avoid contact with eyes, skin, and clothing.

CLEANING OF SPRAYER

Flush with clean water all parts of the sprayer including the pump, hoses and nozzles several times prior to and after the use of this product.

CONDITIONS OF SALE

Seller warrants this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions and use. No one other than an officer of the seller is authorized to make any warranty, guarantee or direction concerning this product. Because time, place, rate of application and other conditions of use are beyond the seller's control, the seller's liability from handling, storage, and use of this product is limited to replacement of the product or a refund of the purchase price. It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions, abnormal conditions, presence of other materials, the manner of application, or other factors, all of which are beyond the control of Valagro USA or the seller. All such risks shall be assumed by buyer.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

NET WEIGHT 5 Lb (2.3 kg)



DO NOT SELL UNPACKED

BREXIL is a Registered Trademark of VALAGRO S.p.A.

Produced and packed by **VALAGRO S.p.A.**

Zona Ind.le - 66041 Ateessa (CH) - ITALY

Distributed and guaranteed by **Valagro USA Inc.**

19500 Hwy 249, suite 245 - Houston TX 77070

Tel (281) 664 8700 - Fax (281) 664 8701



4143374E07F159